Prescribing & Monitoring Training using Plasma Lactate

By
Dr Michael Ponchard

Aims of Presentation

 Demonstrate how the art & science of training can be combined

- Promote the use of sports science methodologies
- Explain how owners, trainers & horses benefit from measuring the performance variable called plasma lactate

Introduction

PhD Program

Mr Fred Kersley

Australian swimming team

Physiological information

• The equine industry

Aim of Training

To win races

Correct ratio of aerobic & anaerobic exercise


Training for 3200 m Vs 1600 m race

Clasification of various speeds

• Snapshot of current fitness status

Energy Systems

- Anaerobic Systems
 - 1. Phosphagen system
 - 2. Lactic acid system
- Aerobic System
 - 3. Aerobic system




Race Model

Start

- Phosphagen system

• Middle phase - Aerobic system



• Last 800m

- Lactic acid system (production)

Finish

- Lactic acid system (tolerance)


Recovery


- Aerobic system (clearance)


Standardbred Races


- Aerobic energy contribution underestimated
- Higher and faster rate of increase in VO₂max
- Aerobic energy probably contributes greater than 70% of race energy











Periodisation of Training

- Divide training program into smaller phases
- Each phase has specific objectives
- Specific types of training & associated lactate concentrations

Phases of training

General preparatory: 8-weeks

• Specific preparatory: 4-weeks


• Pre-competitive: 4-weeks (trials)

• Competitive: 3-6months

• Transition : spelling


Training Terminology

	Types of training	Lactate	Workouts
	Recovery training	1-2 mM	1x20-min
GP	Aerobic threshold	2 mM	2x10-min
	VLa4	4 mM	3x5-min
	Aerobic overload	6-8 mM	3x4-min
SP	Maximal oxygen uptake	10-12 mM	3x3-min
	Lactate tolerance	14-17 mM	2x2-min
PC	Lactate production	20-25 mM	2x1-min
	Sprint training	3-4 mM	3x30 sec




East German Protocol

Each parameter reflected a different aspect of conditioning




Lactate and Performance in Sport



Conditioning Assessment

SLTP Test Running					
	Runners				
	А	С			
Speed_	<u>Lactate</u>	<u>Lactate</u>			
(km/h)	mmol/l	mmoM			
13.5	2.5	2.3			
14.5	3.1	3.5			
15.5	4.5	5.1			
Anaerobic	7.5	9.2			
Test	ſij	ViL			



Characteristics - Anaerobic Profile

• 45-90 seconds - maximum effort

- Time for 800m & collect blood samples 2 4 6 and 8 minutes post-exercise
- The faster the time, or the higher the maximum lactate concentration, the higher the anaerobic capacity
- Sectional times each 200m speed rating

Performance Classification

Performance Ratings & Classifications		Aerobic system VLa4 Sec/400m	Anaerobic system 800 m Sec
3	Poor	43	61
4	Fair	42	60
5	Average	41	59
6	Good	40	58
7	Very good	39	57
8	Excellent	38	56
9	Superb	37	55

Fitness Ratings

Aerobic/Anaerobic Ratings

As fitness progresses

• 3/3 = 6 Poor / Poor = unfit

• 6/4 = 10 Good aerobic / Fair anaerobic

• 7/6 = 13 Very good aerobic / Good anaerobic

• 8/7 = 15 Excellent aerobic / Very good anaerobic

Conclusions

- 1. Fitness profiling provides a performance assessment of the aerobic & anaerobic energy systems
- Identifies ideal training velocities for various types of training
- 3. Demonstrates how each energy system has improved during the previous 4 weeks of training (Trainability)

4. Identifies which energy system should receive more attention in the next 4 weeks of training.

5. When the total performance score plateaus, an indication of the horse's overall genetic potential is obtained

Developing monthly plasma lactate curves reduces the guesswork associated with equine training

Combining the knowledge of the trainer & exercise physiologist enhances the training process

Acknowledgements

- Mr Fred Kersley
- Harness Racing Council